



Colonoscopy Prep Instructions: MOVIPREP SPLIT

Gastro Virginia

Phone: 571-570-1819 Fax: 571-506-2446 | <http://www.gastrocentersva.com>



FOLLOW THESE INSTRUCTIONS EXACTLY. If you have questions, call the nurse line for your location.

WHAT IS A COLONOSCOPY?

A colonoscopy is an examination of the colon (large intestine) with a flexible tube about the thickness of your pinkie finger, which transmits a live color image onto a television screen. For the procedure to be successful, your colon must be completely clean. If it is not adequately prepared, the procedure will need to be rescheduled.

- Provide us with an updated medication list at least 14 days prior to your procedure.
- Pick up your **MOVIPREP prescription** at your pharmacy. Do not wait until the day before.

1 WEEK BEFORE

- Obtain any medical clearances prior to scheduling procedure, if applicable
- Stop **NSAIDs and Iron** 5 days before: Ibuprofen (Advil, Motrin), Naproxen (Aleve, Naprosyn), Aspirin (Bayer, Ecotrin, Miniprin), Diclofenac (Voltaren, Cambia), Celecoxib (Celebrex), Meloxicam (Mobic), Indomethacin (Indocin), Ketorolac (Toradol), and others. **Tylenol (acetaminophen) is OK.**
- Check your medications.** If you are taking **blood thinners** (Coumadin, Warfarin, Eliquis, Brilinta, Clopidogrel, etc.) or **GLP-1/weight loss medications** (Ozempic, Wegovy, Mounjaro, Zepbound, Saxenda, Victoza, Trulicity, Byetta/Bydureon, Adlyxin, Phentermine), **STOP these 7 days before your procedure.**

3 DAYS BEFORE

- Stop all vitamins and supplements. Minimize fiber: Avoid beans, peas, corn, popcorn, nuts, tomatoes, cucumbers, pickles, sesame seeds, strawberries, watermelon, oatmeal, kiwi, grapes, and other foods with small seeds or skins.
- Hold SGLT2 Inhibitors:** Canagliflozin (Invokana), Dapagliflozin (Farxiga), Empagliflozin (Jardiance), Ertugliflozin (Steglatro), Sotagliflozin (Inpefa).

1 DAY BEFORE

MORNING — When you wake up

- Do not eat any food** from the time you wake up. Drink only **clear liquids** all day. See the clear liquids chart below.
- Hold oral GLP-1 medications (e.g., Rybelsus), Metformin & Glyburide today.**
- Prepare solution by mixing **1st pouch A and 1st pouch B** with 1 liter of water then refrigerate.

5:00 PM — Start Dose 1 (Evening Prep)

- Begin drinking the 1 liter of MOVIPREP over one hour (8-ounces every 10 minutes) and then drink ½ liter of clear fluids. If you vomit, wait 30 minutes, then begin drinking again, taking 6-ounces every 15 minutes.
- Mix **2nd pouch A and 2nd pouch B** with 1 liter of water and refrigerate. Continue drinking as many clear liquids as possible after completing the above steps. **Nothing by mouth after midnight** except the second dose of prep.

DAY OF YOUR PROCEDURE

5 Hours Before Your Arrival Time — Start Dose 2

Begin drinking the remaining 1 liter of MOVIPREP preparation solution. Drink 8-ounces every 10 minutes until you finish. If you vomit, wait 30 minutes, then begin drinking again, taking 6-ounces every 15 minutes.

4 Hours Before Your Arrival Time — Nothing More

Do not eat or drink anything 4 hours before your procedure arrival time. No gum, mints, or hard candy. You may brush your teeth but do not swallow.

Take your morning prescribed medications with a sip of water **UNLESS** otherwise directed by your doctor.

If your stool is not clear (see-through, like yellow water) by your procedure morning, call your location's nurse line immediately.

CLEAR LIQUIDS REFERENCE

Clear Liquids — OK TO DRINK	
Black coffee or tea	Water
Clear juice (apple, white grape, white cranberry)	Sports drinks: Gatorade, Powerade, Propel
Clear soda (Sprite, ginger ale, seltzer)	Jell-O, popsicles, Italian ice
Clear broth (chicken, beef, vegetable)	Kool-Aid, Boost Breeze, iced tea, limeade
ABSOLUTELY NO RED OR PURPLE DYE, DAIRY PRODUCTS, ORANGE JUICE, TOMATO JUICE, GRAPEFRUIT JUICE, ALCOHOL, OR ANYTHING WITH A CREAM BASE.	

TRANSPORTATION & CANCELLATION POLICY

Transportation: You cannot drive or operate machinery after your procedure. You must have a friend, family member, or arranged medical transport. Total time in the office is about 2–2½ hours. Your ride does not need to stay — we will contact them 20–30 minutes before discharge. Uber, Lyft, and taxis **are NOT allowed**. Contact our office if you need a medical transportation service.

Cancellations: We understand that from time to time you may need to cancel an appointment for various reasons. Please observe the rules regarding cancellations.

- Any schedule modification requires at least **5 business days' notice** for procedure appointments.
- Procedure cancellations that are made without appropriate notice are subject to a patient charge of **\$200**.
- If you miss your scheduled appointment without appropriate notice, the above charges will be applied accordingly.

NURSE LINE — CONTACT YOUR LOCATION WITH ANY QUESTIONS

Fairfax	(571) 570-1819	Alexandria	(571) 570-1819
---------	----------------	------------	----------------

A nurse will call you within 1 week before your procedure to review prep instructions and answer questions.